

IMPACT OF HYPOTHYROIDISM AND LIPID PROFILE ON OBESITY

Shams UA¹, Zeb MA¹, Karim WA¹, Azhar M¹, ul Haq F² and Muhammad N¹

¹Institute of Paramedical Sciences, Khyber Medical University Peshawar, Pakistan

²University of Health Sciences Lahore Department of Hematology, Pakistan

*Corresponding Author: Shams, Department of Hematology, University of Health Sciences Lahore, Pakistan.

Tel: 00 92 345 9091377, E-mail: isuasbe@gmail.com

HIGHLIGHTS

- The current study aims to determine association of obesity with hypothyroidism, lipid profile and thyroid hormones.
- The higher prevalence of obesity in female is the indicative risk.
- Crystal significant association between obesity and LDL was founded (P-value ≥ 0.005). No association was observed for other study variables.
- Association between obesity and LDL P value 0.004 is the out- come of the study and need an point eye on the food which is the main reason of LDL in food intake of daily life.

Abstract

Aims of the Study: The current study aims to determine association of obesity with hypothyroidism, lipid profile and thyroid hormones.

Methods and Materials: Total of 150 samples collected from obese participants of > 30 Body Mass Index. For thyroid profile samples were run on (cobas e411) principle of the electro generated chemi-luminescence (ECL) and lipid profile on cobas C311 analyze.

Results: Among study participant 104 (69.3%) were females and 46 (30.7%) were males. The higher prevalence of obesity in female is the indicative risk. Result of chi square test predicted; obesity and hypothyroidism p value 0.817,

obesity and triglyceride 0.144, obesity and HDL 0.565, obesity and cholesterol 0.204 and obesity and LDL 0.004.

Conclusion: Crystal significant association between obesity and LDL was founded (P-value ≥ 0.005). No association was observed for other study variables.

Keywords: Obesity; Hypothyroidism; TSH; LDL; HDL; TG; Cholesterol

Citation:

Shams UA, Zeb MA, Karim WA, Azhar M, ul Haq F and Muhammad N: Impact of hypothyroidism and lipid profile on obesity. Asian Journal of allied health sciences. 2023; 8(2):5-13.

Introduction:

Obesity is one of the most salient healths risk of all time. The prevalence of obesity has increased worldwide since the mid-1970s. According to the report of National Health and Nutrition Examination Survey, obesity affected 32.2% of adults in 2003– 2004. Obesity is associated with an increased risk of diabetes, dyslipidemia, kidney disease, cardiovascular disease, all-cause mortality, and cancer.^{1,2}

A BMI of at least 30 is currently generally acknowledged as obesity. Overweight denotes the presence of extra body weight. Obesity denotes the presence of extra deposited fats. Every single obese individual is overweight; however, all overweight people are not obese as

overabundant body weight may emerge from muscles.³ The mechanisms regulating body weight are complex, stimulated by various factors: physiological, societal, environmental, genetic and behavioral. None of those is understood at the individual degree, dietary, metabolic, hormonal and neuronal alerts are incorporated inside the brain to produce modifications in behavior (eating, physical activity) and body metabolism.⁴⁻⁷

Obesity, particularly central obesity, is related to many endocrine abnormalities, such as thyroid disorder.⁸ Thyroid disorder is associated with modifications in body weight and composition, body temperature, and total resting energy expenditure independently of physical hobby. Both subclinical and overt hypothyroidism are often related to weight gain.⁹⁻¹⁰

Hypothyroidism is associated with reduced thermogenesis, decreased metabolic rate, and has additionally been shown to correlate with a higher body mass index (BMI). There is medical evidence suggesting that even slight thyroid disorder is linked to considerable modifications in body weight and represents a hazardous aspect for overweight and obesity.¹¹ An Indian study having sample size of 450 showed that among the obese, 33% had overt, and 11% had subclinical hypothyroidism.¹²⁻¹³ Fat cells active endocrine organs because they produce leptin. The correlation among TSH and BMI might be mediated through leptin. Energy homeostasis is regulated by leptin through informing the brain about fat reserves. Leptin is likewise a crucial neuroendocrine regulator of the hypothalamic-pituitary-thyroid axis through regulation of TRH gene expression within the paraventricular nucleus, and TSH in turn will stimulate leptin secretion through human adipose tissue. Leptin additionally influences thyroid deiodinase activities with activation of T4 to T3.¹⁴

Extreme obesity is related to thyroid disorder because of HPTA anomaly inflicting elevated TSH. Body weight and serum TSH are linked by signals from adipose tissues. TRH is immediately released in the paraventricular nucleus of neurons by stimulation of leptin produced by adipose tissues thus increasing TSH level.¹⁴

The negative feedback among TSH and TRH is decreased in obesity due to reduced T3 receptors [16-18]. Thyroid hormones drastically affects lipoprotein metabolism and risk factors of CVD, thus enhancing overall CVD risk.^{19,20} Even with the slight increase in TSH levels a linear increase in Cholesterol, LDL, TG, and decrease in HDL have been observed.²¹

The literature is still deficient that either a correlation exists among hypothyroidism and obesity. Further, there is clear study conducted yet to find association between lipid profile and thyroid hormones in obese patients. Therefore, this study is conducted to determine the association of obesity with hypothyroidism and lipid profile.

2. Materials and Methods

2.1 Study design and Study sites

Cross-sectional Analytical study was conducted in Khyber Medical University Peshawar. A total of 150 patients with >30BMI and no CVD Patients were included in the study. The study was carried out in Multi Healthcare centre (Lady Reading Hospital, Khyber Teaching Hospital and Hayatabad Medical Complex) in Peshawar.

The study was approved from the institutional review board of Institute of Paramedical Sciences, Khyber Medical University.

2.2 Sampling Technique and procedure

Blood samples were collected in a Yellow top Gel Tube (Clot activator) from B.D Company from the patients of multi health care centres Peshawar

through convenient sampling. The samples were labelled with medical record numbers of concern hospitals. Serum Samples of thyroid profile were run on COBAS e-411 of electro generated chemiluminescence (ECL) principle and for lipid profile samples were run on the COBAS c-311 of photometric assays and ion-selective electrode measurements principle and uses serum/ plasma.

The data was analysed using SPSS. For descriptive statistics cross tabulation was done between different variables and for inferential statistics **Chi-square** test was performed.

3. Results:

150 Obese participants were screened for key variables i.e. thyroid profile & lipid profile. These results showed the high number of female obese participant which may be due to their home life style of that particular regions.

3.1 Age and Gender wise distribution of Participants

150 patients having 104 (69.3%) females and 46 (30.7%) males. The most prevalent age group is 31-41 years which need further assessment. Age groups 10-20 years were the more prevalent group among obese study participants.

Following Table 1 and Figure 1 shows the frequency and percentage of the said variables.

Variables	Frequency (%)
Age	
10-20	31 (20.7%)
21-30	20 (13.3%)
31-40	38 (25.3%)
41-50	27 (18.0%)
51-60	19 (12.7%)
61-70	12 (8.0%)
71-80	2 (1.3%)
81-90	1 (0.7%)

Sex	
Male	46 (30.7%)
Female	104 (69.3%)
Weight	
60-70	6 (4%)
71-80	41 (27.3%)
81-90	70 (46.7%)
91-100	31 (20.7%)
101-110	2 (1.3%)
Obese Class	
Over-Weight	4 (2.7%)
Obese Class I	136 (90.7%)
Obese Class II	10 (6.7%)

Table 1: Baseline characteristic of study participants (n=150)




Figure 1: Age wise distribution of study participants




Figure 2: Gender wise distribution of study participants

3.2 Weight wise distribution

71-80 kg was observed among 41 (27.3%) study subjects as well as 81-90 kg among 70(46.7%). The lowest Weight of obesity is 60-70 which indicate alternative reason that small height participant have low chance of obesity. Following Table 1 and Figure 3 shows the frequency and percentage of said variables.




Figure 3: Weight difference among study participants

3.3 Obese class distribution

Obese class 1 is the main obesity found in our study 136 (90.7%) while other classes of obesity have low prevalence (Table 1 and Figure 4).




Figure 4: Classification of obese participants

3.3.1 Obese class and fast food consumption

It was seen that the fast food consumption among obese classes was high in class 1 obese which is the man age where people want go outside and eat as shown in Table 2a.

Obese Classes	Fast Food					Total
	Zinger	Pizza	Chinese soup	Others	No	
Over-Weight	1	0	0	0	3	4
Obese Class I	24	20	6	1	85	136
Obese Class II	1	0	0	0	9	10
Total	26	20	6	1	97	150

Table 2a: Use of fast food by different obese participants

3.3.2 Obese class and Diet

It was seen that both Veg and Non-Veg was the major meal plan among obese classes. But these results are against the natural and need to be verifying by the comparison of other food in relevant to vegan (Table 3).

Obese Classes	Diet			Total
	Veg	Non-Veg	Both	
Over-Weight	1	0	3	4
Obese Class I	12	3	121	136
Obese Class II	0	0	10	10
Total	13	3	134	150

Table 3: Diet use among different obese class participants

3.3.2 Obese class and Diet

It was seen that both Veg and Non-Veg was the major meal plan among obese classes. But these results are against the natural and need to be verifying by the comparison of other food in relevant to vegan (Table 3).

ObeseClasses	Diet			Total
	Veg	Non -Veg	Both	
Over -Weight	1	0	3	4
ObeseClass I	12	3	121	136
ObeseClass II	0	0	10	10
Total	13	3	134	150

Table 3: Diet use among different obese class participants

3.3.3 Obese class and rice consumption

It was seen that rice consumption have association with obesity. So proper intake of rice should be preferred in obese person Table 2b.

ObeseClasses	Riceconsumption					Total
	1-2timesa week	2-3timesa week	4-5timesa week	4-8timesa month	Nb	
Over -Weight	1	1	1	0	1	4
ObeseClass I	47	47	14	7	21	136
ObeseClass II	5	3	1	0	1	10
Total	53	51	16	7	23	150

Table 2b: Rice consumption by different obese class participants

3.3.4 Obese class and Routine daily

Mild exercise was observed among obese classes. Mild exercise is not enough to overcome the obesity so you need to be more active to be non-obese (Table 4).

ObeseClasses	Routinedailyexercise				Total
	walk/ running	home workouts	strenuous activities	Nb	
Over -Weight	0	1	1	2	4
ObeseClass I	35	18	3	78	136
ObeseClass II	1	2	0	7	10
Total	37	21	4	87	150

Table 4: Routine daily exercise by different obese class participant

3.3.5 Obese class and Other Complication:

Arthritis, back pain, and other complications such as weak eye sight, generalized weakness, were seen among obese classes which indicate obesity is related with other health risk factor and outcome (Table 5).

Obese Classes	Othercomplications					Total
	Frequent infection	arthritis	backpain	others	Nb	
Over -Weight	0	2	0	0	2	4
ObeseClass I	10	21	27	17	61	136
ObeseClass II	1	5	2	0	2	10
Total	11	28	29	17	65	150

Table 5: Complication associated with obese class participants

3.3.6 Obese class and Education

It was observed that most of the obese participants were illiterate which indicate our society need obesity related social work about their harmful impact on health (Table 6).

Obese Classes	Education					Total
	Illiterate	SSC	F. Sc	Bachelor	M.Phil/ PhD	
Over -Weight	3	1	0	0	0	4
Obese Class I	77	23	24	8	4	136
Obese Class II	8	1	1	0	0	10
Total	88	25	25	8	4	150

Table 6: Education level of obese class participants

3.3.6 Obese class and Hypothyroidism

It was observed in our study that there was no association between obesity and hypothyroidism, P value 0.817 (Table 7).

Obese Classes	Hypothyroidism		Total	P.Value
	Yes	No		
Over -Weight	0	4	4	.817
Obese Class I	11	125	136	
Obese Class II	1	9	10	
Total	12	138	150	

Table 7: Hypothyroidism among obese class participants

3.3.6 Obese class and Triglyceride

It was observed in our study that there was no association between obesity and Triglyceride P value 0.144 (Table 8).

Obese Classes	Triglyceride			Total	P. Value
	100 -150	151 -200	201 -250		
Over -Weight	0	4	0	4	.144
Obese Class I	17	91	28	136	
Obese Class II	1	4	5	10	
Total	18	99	33	150	

Table 8: Triglyceride among obese class participants

3.3.7 Obese class and Cholesterol:

It was observed in our study that there was no association between obesity and Cholesterol P value 0.204 (Table 9).

Obese Classes	Cholesterol			Total	P. Value
	151 -200	201 -250	251 -300		
Over -Weight	0	2	2	4	.204
Obese Class I	11	76	49	136	
Obese Class II	3	4	3	10	
Total	14	82	54	150	

Table 9: Cholesterol level among obese class participants

3.3.8 Obese class and LDL

It was observed in our study that there was an clear association between obesity and LDL. P value 0.004. There was an increase in serum LDL among obese classes (Table 10).

Obese Classes	LDL			Total	P. Value
	50-100	101-150	151-200		
Over-Weight	3	0	1	4	0.004
Obese Class I	100	34	2	136	
Obese Class II	10	0	0	10	
Total	113	34	3	150	

Table 10: LDL level among obese class participants

3.3.9 Obese class and HDL

It was observed in our study that there was no association between obesity and HDL. P value 0.565 (Table 11).

Obese Classes	HDL				Total	P. Value
	20-30	31-50	51-70	71-90		
Over -Weight	1	3	0	0	4	.565
Obese Class I	16	68	46	6	136	
Obese Class II	1	3	5	1	10	
Total	18	74	51	7	150	

Table 11: HDL level among obese class participants

4. Discussions

Very limited literature exists on the association of obesity with hypothyroidism and lipid profile, especially in Pakistan and specifically in Khyber Pakhtunkhwa, therefore this study is aimed to determine the association between obesity and hypothyroidism with lipid profile. There was a clear association between obesity and LDL P value 0.004, as LDL was increased in participants. In our study it was observed that there was no association between obesity and hypothyroidism P value 0.817 furthermore it was observed there was no association between obesity and triglyceride, HDL, and cholesterol, P value 0.144, 0.565, 0.204. The higher prevalence of obesity in females is indicative risk and reason behind their obesity. In a study by Abhyuday Verma, MD, Muthu Krishnan Jayaraman it was shown that there is an association between obesity and hypothyroidism p value 0.02. Their sample size was 1075 individuals and had a control group of 450 individuals. In their study the group 1 had 625 individuals in which 44% had BMI of >25. They observed that in obese individuals there was a high prevalence of hypothyroidism 33% and 11% had subclinical hypothyroidism. Detail studies are required to assess the cause and effect relationship between obesity and hypothyroidism.¹³

In a study by Okan et al, it was shown that there was no association between obesity and hypothyroidism. Their sample size was 947 and had a control group of 50 individuals.²²⁻²⁵

They had 4 groups which were based on circumference and BMI and showed a positive relation between BMI, circumference and body fat percentage P value 0.001.²⁶⁻²⁸ They compared all the 4 groups with the control group for serum TSH levels and found no significance in the control group and the other 4 groups P value

0.34, 0.69, 0.12 and 0.47 respectively.²⁹⁻³⁰

In a study by Debmalya Sanyal and Moutusi Raychaudhuri showed no clear association between obesity and Hypothyroidism and stated that further research is necessary to determine whether hypothyroidism is one of the cause of obesity.¹² The difference in results of our study and other study is because most of the studies had a wide sample size average 900+ as compared to our study it was 150 individuals. Most importantly these studies had a control group where as in our study there was no control group due to limited resources. These studies were conducted mostly in the Endocrinology Department of Hospitals therefore they had referral biasness in their results. These studies had advance equipment's that can detect free thyroid hormones while we had no such facilities to do so. These studies were conducted for a long duration such as 2-3 years while our study was conducted for 2 months only. These studies were mostly retrospective while our study was cross sectional.³¹⁻³⁵

5. Conclusion

Association between obesity and LDL P value 0.004 is the out-come of the study and need an point eye on the food which is the main reason of LDL in food intake of daily life.

REFERENCES:

- 01- World Health Organization WHO (2000) Obesity: preventing and managing the global epidemic, Geneva, Sweden.
- 02- Golden SH, Robinson KA, Saldanha I, Anton B, Ladenson PW, JT JoCE (2009) Metabolism. Prevalence and incidence of endocrine and metabolic disorders in the United States: a comprehensive review 94: 1853-78.

- 03- World Health Organization WHO (2000) The Asia-Pacific perspective: redefining obesity and its treatment: Sydney: Health Communications Australia.
- 04- World Health Organization WHO (1998) Obesity: preventing and managing the global epidemic: report of a WHO consultation on obesity, Geneva, Sweden.
- 05- Kumanyika SK (1993) Special issues regarding obesity in minority populations 119:650-4.
- 06- Hodge AM, Zimmet PZJB (1994) Metabolism In: The epidemiology of obesity 8: 577-99.
- 07- Nanan DJJ (2002) The obesity pandemic-implications for Pakistan 52: 6-11.
- 08- Biondi B (2010) Thyroid and obesity: an intriguing relationship, Oxford University Press, UK.
- 09- Hoogwerf BJ, Nuttall FQJTA (1984) Long-term weight regulation in treated hyperthyroid and hypothyroid subjects 76: 963-70.
- 10- A'svold BO, Bjørro T, Vatten LJ (2009) Metabolism. Association of serum TSH with high body mass differs between smokers and never-smokers 94: 5023-7.
- 11- Danforth E, Horton E, O'Connell M, Sims E, Burger A, et al. (1979) Dietary-induced alterations in thyroid hormone metabolism during overnutrition 64: 1336-47.
- 12- Sanyal D, Raychaudhuri MJ (2016) Metabolism Hypothyroidism and obesity: An intriguing link 20: 554.
- 13- Rosenbaum M, Hirsch J, Murphy E, Leibel RL (2000) Effects of changes in body weight on carbohydrate metabolism, catecholamine excretion, and thyroid function 71: 1421-32.
- 14- Verma A, Jayaraman M, Kumar HK, Modi KD (2008) Hypothyroidism and obesity. Cause or effect? 29: 1135-8.
- 15- Thompson GR, Soutar AK, Spengel FA, Jadhav A, Gavigan S, et al. (1981) Defects of receptor-mediated low density lipoprotein catabolism in homozygous familial hypercholesterolemia and hypothyroidism in vivo. 78: 2591-5.
- 16- Seppel T, Kosel A, Schlaghecke RJ (1997) Bioelectrical impedance assessment of body composition in thyroid disease 136: 493-8.
- 17- Pinkney JH, Goodrick SJ, Katz J, Johnson AB, Lightman SL, et al. (1998) Leptin and the pituitary-thyroid axis: a comparative study in lean, obese, hypothyroid and hyperthyroid subjects 49: 583-8.
- 18- Wolf M, Weigert A, Kreymann GJ (1996) Body composition and energy expenditure in thyroidectomized patients during short-term hypothyroidism and thyrotropin-suppressive thyroxine therapy 134: 168-73.
- 19- Burman KD, Latham KR, Djuh Y-Y, Smallridge RC, Tseng Y-CL, et al. (1980) Solubilized nuclear thyroid hormone receptors in circulating human mononuclear cells 51: 106-16.
- 20- Mihály E, Fekete C, Tatro JB, Liposits Z, Stopa EG, et al. (2000) Hypophysiologic thyrotropin-releasing hormone- synthe-

- sizing neurons in the human hypo-thalamus are innervated by neuropeptide Y, agouti-related protein, and α -melanocyte-stimulating hormone 85: 2596-603.
- 21- Blum WF, Englaro P, Attanasio AM, Kiess W, Rascher WJ (1998) Human and clinical perspectives on leptin 57: 477-85.
- 22- Tangkilisan AH, Akune KJPI (2007) Some factors related to lipid profile in obese children at junior high schools in Manado 47: 166-71.
- 23- Teixeira PJ, Sardinha LB, Going SB, Lohman TG (2001) Total and regional fat and serum cardiovascular disease risk factors in lean and obese children and adolescents 9: 432-42.
- 24- Duntas LHJT (2002) Thyroid disease and lipids 12: 287-93.
- 25- Canaris GJ, Manowitz NR, Mayor G, Ridgway EC (2000) The Colorado thyroid disease prevalence study. 160: 526-34.
- 26- Åsvold BO, Vatten LJ, Nilsen TI, Bj̊ro TJ (2007) The association between TSH within the reference range and serum lipid concentrations in a population-based study. The HUNT Study 156: 181-6.
- 27- Bakker O, Hudig F, Meijssen S, Wiersinga WMJB (1998) Effects of triiodothyronine and amiodarone on the promoter of the human LDL receptor gene 249: 517-21.
- 28- Shin D-J, Osborne TF (2003) Thyroid hormone regulation and cholesterol metabolism are connected through sterol regulatory element-binding protein-2 (SREBP-2) 278: 34114-8.
- 29- Faure P, Oziol L, Artur Y, Chomard PJB (2004) Thyroid hormone (T3) and its acetic derivative (TA3) protect low-density lipoproteins from oxidation by different mechanisms 86: 411-8.
- 30- Iglesias P, Díez JJJC (2007) Influence of thyroid dysfunction on serum concentrations of adipocytokines. 40: 61-70.
- 31- Rizos C, Elisaf M, Liberopoulos EJ (2011) Effects of thyroid dysfunction on lipid profile 5: 76.
- 32- Richter MM (2004) Electrochemiluminescence (ecl) 104: 3003-36.
- 33- Diagnostics Roche (2001) CRcT Cobas e 411 analyser, Ltd, USA.
- 34- Tansley B (1998) Photometric measurement apparatus, Google Patents, USA.
- 35- Bakiner O, Bozkirli E, Cavlak G, Ozsahin K, Ertorer EJ (2014) Are plasma thyroid-stimulating hormone levels associated with degree of obesity and metabolic syndrome in euthyroid obese patients? A Turkish cohort study