

Association Between Patient's Characteristics and Hamstring Flexibility in Patients with Knee Osteoarthritis

Komal Tariq^{1*}, Filza Shoukat¹, Umair Ahmed¹, Farwa Batool¹ and Aruj Fatima¹

¹University Institute of Physical Therapy, Faculty of Allied Health Sciences, University of Lahore, Lahore, Pakistan.

*komaljutt358@gmail.com

Highlights:

- Osteoarthritis is most common skeletal disease.
- Patients characteristics such as occupation have association with hamstring muscle in osteoarthritis

Abstract:

Background: Knee osteoarthritis is most common among patients. Patients with knee Osteoarthritis usually have pain, brittle bone and felt difficulty during movement. Mostly individuals after forty years of age have felt the symptom of knee osteoarthritis.

Objective:

The objective of this study was to find out the association between patients characteristic and hamstring flexibility in knee osteoarthritis.

Methodology:

A cross sectional study was conducted in one hundred and one patients that were selected from rehabilitation centre of Allied hospital, Faisalabad. The study duration was six months. Participants with knee osteoarthritis (stage 1 & 2), age forty to sixty five years were involved in this study if they had nerve related issues, bone or muscle related disorder were not counted in this study. Subject's characteristics such as their profession, age, pain location as well as duration of pain related questions were asked from patients. Active knee extension test was used for recording the hamstring length. Chi square test was used to determine the association.

Results:

Men were 38(37.6%) and women 63(62.4%).

Occupations of patients had influence on hamstring flexibility ($P<0.000$) while patient's age ($P<0.071$) durations of pain ($P<0.856$) and pain region ($P<0.956$) had not associated with hamstring flexibility.

Conclusions:

It is concluded that patient's characteristic in which only occupations of patients had compromised the hamstring flexibility in knee osteoarthritis subjects rather than age, duration of pain and pain region.

Keywords:

Patient's characteristics, hamstring flexibility, Osteoarthritis

Introduction:

Knee osteoarthritis is most common among patients. Patients with knee Osteoarthritis usually have pain, brittle bone and felt difficulty during movement. Majority of individuals after forty years of age have felt the symptom of knee osteoarthritis. Approximately 3.8% of the people around the globe having the knee osteoarthritis which is equalize to two hundred and seventy seven individuals that have been experience the knee osteoarthritis.¹⁻⁴ Individuals with overweight also have linked with knee osteoarthritis. Body over weight affects the knee mechanic or exerts pressure on joint and individuals have complained of arthritis. Females are more suffered from Knee osteoarthritis as compared to males due to overweight issues.⁵ Some studies stated that higher blood pressure and glucose level also affects the knee joint as well as subject with this issue have complained to knee osteoarthritis.^{6,7} Knee osteoarthritis cause pain and

functional disability worldwide.⁸ Osteoarthritis frequency increases with the passage of time so strategies are necessary to be taken to reduce the impact of knee osteoarthritis on patients through primary and secondary prevention.⁹ Osteoarthritis is most common between middle aged populations whose reduced quality of life and loss to productivity leads to high disease cost.¹⁰ Personal and economic factor cause widespread pain, but its etiology is not exact known. Longitudinal studies describe that anxiety, cognitive decline, poor sleep, and multi joint osteoarthritis, poor health-related quality of life (QOL) baseline pain status, older age and a family history of chronic pain all are the risk factor for widespread pain. However, both non modifiable and modifiable risk factors have been recognized but exact mechanisms through which these risk factors may lead to widespread pain have not been investigated.¹¹ A cross-sectional study of patients with knee osteoarthritis show a relation among high pain severity and low-grade radiographic knee Osteoarthritis¹², declared that may be pain itself rather than structural pathology may lead to widespread pain. Hamstring muscle played important role in maintenance of knee joint function during walking or any physical activity. Hamstrings muscle produced the hip extension and knee flexion there are many factors that cause or linked with knee osteoarthritis. But the reason to conducted this study have to evaluated the subjects demographic characteristic either have linked with or not to hamstring flexibility in knee osteoarthritis because subjects occupational level and activity level both have impact on knee joint or muscle activity and disturbed their knee joint mechanic and articular joint as well as bone. So, the purpose of this study was to evaluate either the subject's characteristic have association with knee

osteoarthritis or not.

Methodology:

A descriptive cross sectional study was conducted. Patients' information was obtained from rehabilitation centre of Allied hospital Faisalabad after taking the ethical approval letter. One hundred and one patients were participated in this study. Permission forms were signed from all patients to take part in this study. Participants with knee osteoarthritis (stage 1 & 2), age forty to sixty five years were involved in this study if they had nerve related issues, bone or muscle related disorder were not counted in this study. AKT (active knee extension test) was used to evaluate the hamstring muscle length.¹³ Participant's demographical data were obtained by asking questions about their duration of pain, their occupation, location of pain and age. Chi square test was used to find out the association between participants characteristic with hamstring muscle flexibility

Results:

Out of 101 (100%) males were 38(37.6%) and females 63(62.4%). Majority of the respondents 27.7 % (28) were belongs to 40 to 45 years age group. Chi square test showed that there was no association between age of respondents and hamstring flexibility in knee osteoarthritis ($P < 0.071$) (Table 1). Majority of respondents 42(41.6%) had pain in patellofemoral joint. So, there was no association between pain region and hamstring flexibility in knee osteoarthritis ($P < 0.965$) (Table 2).

Age	Frequency(%)	P-Value
40 - 45	28 (27.7%)	0.071
46 - 50	21 (20.8%)	
51 - 55	17 (16.8%)	
56 - 60	16 (15.87%)	
61 - 65	19 (18.8%)	

Table 1: Descriptive statistics

Variable	Frequency(%)	P-Value
Pain Region		
Patellofemoral Joint Pain	42 (41.6%)	0.956
Tibiofemoral Joint Pain	23 (22.8%)	
Bilateral Joint Pain	36 (35.6%)	

Table 2: Association of age and pain region with hamstring flexibility

Mostly respondents were females, and they were housewives. Housewives 52(51.5%) were mostly facing the knee problems. Secondly there were workers and thirdly businessmen who were facing the knee problems. They both were 10(9.9%) and 9(8.9%) of the total respondents. These all individuals had hamstring tightness so, there was association of occupations with hamstring flexibility in knee osteoarthritis ($P<0.000$) (Table 3)

Variable	Frequency (%)	Variable	Frequency (%)	P-Value
Businessman	9 (8.9%)	Policeman	1 (1.0%)	0.000
designer	1 (1.0%)	Radiologist	1 (1.0%)	
Driver	2 (2.0%)	Retired	1 (1.0%)	
Electrician	1 (1.0%)	Retired	1 (1.0%)	
House Wife	52 (51.5%)	Officer	3 (3.0%)	
Jeweler	1 (1.0%)	Security Guard	6 (5.9%)	
Lawyer	1 (1.0%)	Servant	2 (2.0%)	
Milkman	1 (1.0%)	Tailor	6 (5.9%)	
Nurse	1 (1.0%)	Teacher	10 (9.9%)	
Peon	1 (1.0%)	Worker		

Table 3: Descriptive statistics and association between occupation and hamstring flexibility

Many of the respondents 20.8 % (21) were facing the knee problems for last 2 months. 14.9 % (15) of the respondents were facing the knee problem since last one month. 12.9 % (13) of the respondents were facing the knee problem from last 3 months. So there was association between duration of pain and hamstring flexibility in osteoarthritis ($P<0.856$).

Duration of Pain	Frequency (%)	Variable	Frequency (%)	P-Value
2 Days	1 (1.0%)	3 Months	13 (12.9%)	0.856
10 Days	1 (1.0%)	3.5 Months	1 (1.0%)	
15 Days	2 (2.0%)	4 Months	3 (3.0%)	
1 Week	3 (3.0%)	5 Months	5 (5.0%)	
2 Weeks	7 (6.9%)	6 Months	1 (1.0%)	
3 Weeks	4 (4.0%)	8 Months	7 (6.9%)	
6 Weeks	5 (5.0%)	9 Months	2 (2.0%)	
8 Weeks	2 (2.0%)	1 Years	1 (1.0%)	
1 Month	15 (14.9%)	2 Years	1 (1.0%)	
1.5 Months	2 (2.0%)	5 Years	1 (1.0%)	
2 Months	21 (20.8%)	15 Years	2 (1.0%)	
2.5 Months	1 (1.0%)		1 (1.0%)	

Table 4: Descriptive statistics and Association between pain duration and hamstring flexibility

Discussion:

This study reported that subject's characteristics such as occupation and duration of pain had compromised the hamstring flexibility while the pain regions and age had not compromised the hamstring flexibility in knee osteoarthritis patients. It means patients occupation and duration of pain had associated with hamstring flexibility.

Fitzgerald *et al.*, claimed that subject's physical feature (muscle power) or outward form, lineament or non-physical or intellectual factor, emotional states had also linked with knee Osteoarthritis disease. If this entire element were changed in subjects their pain sensation and activity level compromised with passage of time. All subjects' information was getting by WOMAC questionnaire (Western Ontario and McMaster Universities Osteoarthritis Index).¹⁴ This present study also reported the result that activity level compromised during occupation had influence on hamstring extensibility.

Tharakan *et al.*, explained that tibiofemoral degradation had compromised the strength of hamstring muscle. They had also compromised the function of knee joint ($P<0.001$) in knee Osteoarthritis patients as well as their pattern of

walking. They had evaluated the muscle flexibility by PKET (passive knee extension test. This study results showed that tibiofemoral joint pain had not compromised the hamstring extensibility in knee Osteoarthritis patients. But duration of pain affects the knee mechanic and flexibility of hamstring muscle ($P<0.000$).¹⁵ But this present study also showed that tibiofemoral joint pain had not any effect on hamstring length ($P<0.956$). But duration of pain had not support the previous study because it had not compromised the hamstring flexibility ($p=0.865$).

Abolahrari *et al.*, proclaim that muscles that are inserted at the knee joint had been affected in knee arthritis patients. This study was done on 23 normal females. VAS was used for pain. This study concluded that muscle lengths were compromised in this disease patient due to pain ($p<0.0001$)¹⁶ and duration of pain also affected the muscle flexibility especially of hamstrings ($p<0.0001$). But in this latest study a hamstring muscles were also affected ($p<0.000$) while duration of pain had not affected the hamstring muscle group in knee Osteoarthritis patients.

King *et al.*, showed that overweight or obesity also had relation with knee Osteoarthritis disease because over weight affect the joint and their load shift result in joint dreadful conditions, lead to continuance of this disease ($P=0.039$). More muscle mass progress the cartilage obliteration as well as inner bone mass¹⁷ if overweight is reduced Osteoarthritis subject relived from pain.^{18,19} But this study had not told about the overweight or obesity either compromise the flexibility or not.

A study of Bednarz *et al.*, result showed that occupation had effect on knee joint mechanic as well as hamstring muscle length. Different functional activities while doing their work affect muscle flexibility ($p<0.002$).²⁰ This present study results also

support this study because occupations of subjects had relation with hamstring muscle flexibility ($p<0.000$)

Conclusions:

In case of knee osteoarthritis, patient's characteristic in which only occupations of patients had association with hamstring flexibility rather than age, location of pain and pain region. It is recommended to conduct a research to find out the association of BMI with hamstring flexibility in osteoarthritis patients with large sample size.

References:

1. Cross M, Smith E, Hoy D, Nolte S, Ackerman I, Fransen M, *et al.*, The global burden of hip and knee osteoarthritis: estimates from the global burden of disease 2010 study. *Annals of the rheumatic diseases*. 2014 Jul 1;73(7):1323-30..
2. U.S. Census Bureau. International data base [Internet]. Washington, DC: U.S. Census Bureau; 2016.
3. Sancheti P, Shetty VD, Dhillon MS, Sprague SA, Bhandari M. India-based knee osteoarthritis evaluation (iKare): A multi-centre cross-sectional study on the management of knee pain and early osteoarthritis in India. *Clinics in orthopedic surgery*. 2017 Sep 1;9(3):286-94.
4. Eymard F, Parsons C, Edwards MH, Petit-Dop F, Reginster JY, Bruyère O, *et al.*, Diabetes is a risk factor for knee osteoarthritis progression. *Osteoarthritis and cartilage*. 2015 Jun 1;23(6):851-9.
5. Yusuf E, Nelissen RG, Ioan-Facsinay A, Stojanovic-Susulic V, DeGroot J, van Osch G, Association between weight or body mass index and hand osteoarthritis: a systematic review. *Annals of the rheumatic diseases*. 2010 Apr

- 1;69(4):761-5.
6. Zhang YM, Wang J, Liu XG. Association between hypertension and risk of knee osteoarthritis: A meta-analysis of observational studies. *Medicine*. 2017 Aug;96(32).
7. Maeda Y, Nakamura N, Sugano N. Perioperative hyperglycemia and postoperative periprosthetic joint infection (PJI) after total knee and hip arthroplasty. *Ann Musculoskelet Med*. 2018;2(1):001-5.
8. Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, Ezzati M, *et al.*, Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *The Lancet*. 2012 Dec 15;380(9859):2163-96.
9. Cross M, Smith E, Hoy D, Nolte S, Ackerman I, Fransen M, *et al.*, The global burden of hip and knee osteoarthritis: estimates from the global burden of disease 2010 study. *Annals of the rheumatic diseases*. 2014 Jul;73(7):1323-30.
10. Losina E, Paltiel AD, Weinstein AM, Yelin E, Hunter DJ, Chen SP, *et al.*, Lifetime medical costs of knee osteoarthritis management in the United States: impact of extending indications for total knee arthroplasty. *Arthritis care & research*. 2015 Feb;67(2):203-15.
11. McBeth J, Lacey RJ, Wilkie R. Predictors of new-onset widespread pain in older adults: results from a population-based prospective cohort study in the UK. *Arthritis & Rheumatology*. 2014 Mar;66(3):757-67.
12. Riddle DL, Stratford PW. Knee pain during daily tasks, knee osteoarthritis severity, and widespread pain. *Physical therapy*. 2014 April 1;94(4):490-8.
13. Shamsi M, Mirzaei M, Khabiri SS. Universal goniometer and electro-goniometer intra-examiner reliability in measuring the knee range of motion during active knee extension test in patients with chronic low back pain with short hamstring muscle. *BMC Sports Science, Medicine and Rehabilitation*. 2019 Dec;11(1):4.
14. Tharakan MV, Gopalswami AD. A Study to Determine the Influence of Tibiofemoral Degeneration over Hamstring Flexibility and Functional Performance. *Indian Journal of Physiotherapy and Occupational Therapy-An International Journal*. 2016 Jul;10(3):102-7.
15. Fitzgerald GK, White DK, Piva SR. Associations for change in physical and psychological factors and treatment response following exercise in knee osteoarthritis: an exploratory study. *Arthritis care & research*. 2012 Nov;64(11):1673-80.
16. Abolahrari Shirazi S, Ghafari Nezhad F, Ebrahimian M, Nouraddini E, Mansoorian A, Emami F. Flexibility of knee Joint muscles in women with knee osteoarthritis and healthy controls. *Journal of Rehabilitation Sciences & Research*. 2016 Mar 21;2(3):47-52.
17. Berry PA, Wluka AE, Davies-Tuck ML, Wang Y, Strauss BJ, Dixon JB, *et al.*, The relationship between body composition and structural changes at the knee. *Rheumatology*. 2010 Aug 18;49(12):2362-9.
18. King LK, March L, Anandacoomarasamy A. Obesity & osteoarthritis. *The Indian journal of medical research*. 2013 Aug;138(2):185.
19. Thijssen E, van Caam A, van der Kraan PM. Obesity and osteoarthritis, more than just wear and tear: pivotal roles for inflamed adipose

tissue and dyslipidaemia in obesity-induced osteoarthritis. *Rheumatology*. 2014 Dec 11;54(4):588-600.

20. Bednarz J, Kolatorowicz A, Lowe CM. Activity level influences the likelihood of developing knee osteoarthritis. *Osteoarthritis and Cartilage*. 2019 Apr 1;27:S279.